

Publicaties der leden 2008

Analytisch: Hematologie

- Stroobants AK, van Oerle R, Berends F, Sturk A, Spronk H, Henskens YMC. Oude technieken in een nieuw jasje. Ned Tijdschr Klin Chem Labgeneesk. 2008; 33: 260-5.
- van den Dool EJ, Henskens YMC, Stroobants AK. Anti-factor-Xa-activiteitbepaling voor verschillende heparines, heparinoïde en pentasaccharide. Ned Tijdschr Klin Chem Labgeneesk. 2008; 33: 244-6.
- de Kok JB, van der Meer J, Stroobants AK, Henskens Y, Ponjee G, van Wijk E, Hackeng C, Schamhorst V, Mulder A. Veneuze trombo-embolie: wanneer zijn laboratoriumtesten voor trombofilie zinvol? Ned Tijdschr Klin Chem Labgeneesk. 2008; 33: 228-32.
- Henskens YMC, Stroobants AK, van den Dool EJ, Hamulyak K, van den Besselaar T. Therapiemonitoring van antistollingsbehandeling. Ned Tijdschr Klin Chem Labgeneesk. 2008; 33: 238-43.
- Stroobants AK, Heckman M, Bakker B, van den Dool EJ, Sturk A. Gevoeligheid van verschillende APTT-reagentia. Ned Tijdschr Klin Chem Labgeneesk. 2008; 33: 250-4.
- Akkerman JW. From low-density lipoprotein to platelet activation. Int J Biochem Cell Biol. 2008; 40 (11): 2374-8.
- Boonstra JG, van Toorenbergen AW. De bepaling van serum vrije lichte ketens. Ned Tijdschr Hematol 2008; 5: 260-5.
- de Vooght KM, van Wijk R, van Wesel AC, van Solinge WW. Characterization of the -148C>T promoter polymorphism in PKLR. Haematologica. 2008 Sep; 93 (9): 1407-8.
- de Vries WB, Karemaker R, Mooy NF, Strengers JL, Kemperman H, Baerts W, Veen S, Visser GH, Heijnen CJ, van Bel F. Cardiovascular follow-up at school age after perinatal glucocorticoid exposure in prematurely born children: perinatal glucocorticoid therapy and cardiovascular follow-up. Arch Pediatr Adolesc Med. 2008 Aug; 162 (8): 738-44.
- Folman CC, Ligthart PC, Smid WM, de Vooght KMK, Adriaansen HJ. Een zwangere vrouw met antistoffen anti-CRAM, gericht tegen een hoogfrequent antigen: serologisch en klinisch beloop. Tijdschrift voor Bloedtransfusie 2008; 1: 73-7.
- Nieuwland R, Stroobants AK, Sturk A. Hemostase anno 2008. Ned Tijdschr Klin Chem Labgeneesk. 2008; 33: 223-8.
- Huisman A, van Solinge WW. Alternatief stroomschema voor laboratoriumonderzoek van anemie (rebuttal). Ned Tijdschr Geneesk. 2008; 52: 586-9.
- Novotny VMJ, van Solinge WW, van Wijk R. Erfelijke hemolytische anemieën. Handboek Hematologie, De Tijdstroom Uitgeverij, Utrecht (2008), hoofdstuk 7, pp 81-99
- Pruijsten RV, de Jonge N, Kirkels JH, Klopping C, Doevendans PA, Oosterom A, Kemperman H, Lahpor JR. Left ventricular assist device: a functional comparison with heart transplantation. Neth Heart J. 2008 Feb; 16 (2): 41-6.
- Raijmakers MTM, Kuijper PHM, Bakkeren DL, Vader HL. The effect of paraproteins on the erythrocyte sedimentation rate: a comparison between the StarrSed and TEST 1. Ann Clin Biochem. 2008 Nov; 45 (Pt 6): 593-7.
- Ringrose JH, van Solinge WW, Mohammed S, O'Flaherty MC, van Wijk R, Heck AJ, Slijper M. Highly efficient depletion strategy for the two most abundant erythrocyte soluble proteins improves proteome coverage dramatically. J Proteome Res. 2008 Jul; 7 (7): 3060-3.
- Thijssen JH. The first 30 years of MATURITAS. Maturitas. 2008 Sep-Oct; 61 (1-2): 1-3.
- Torrance HL, Voorbij HA, Wijnberger LD, van Bel F, Visser GH. Lung maturation in small for gestational age fetuses from pregnancies complicated by placental insufficiency or maternal hypertension. Early Hum Dev. 2008 Jul; 84 (7): 465-9.
- van den Akker EL, Baan CC, van den Berg B, Russcher H, Joosten K, Hokken-Koelega AC, Lamberts SW, Koper JW. Ficoll-separated mononuclear cells from sepsis patients are contaminated with granulocytes. Intensive Care Med. 2008 May; 34 (5): 912-6.
- van Lier M, Verhoef S, Cauwenberghs S, Heemskerk JW, Akkerman JW, Heijnen HF. Role of membrane cholesterol in platelet calcium signalling in response to VWF and collagen under stasis and flow. Thromb Haemost. 2008 Jun; 99 (6): 1068-78.
- van Rijn BB, Franx A, Sikkema JM, van Rijn HJ, Bruinse HW, Voorbij HA. Ischemia modified albumin in normal pregnancy and preeclampsia. Hypertens Pregnancy. 2008; 27 (2): 159-67.
- van Rijn BB, Franx A, Steegers EA, de Groot CJ, Bertina RM, Pasterkamp G, Voorbij HA, Bruinse HW, Roest M. Maternal TLR4 and NOD2 gene variants, pro-inflammatory phenotype and susceptibility to early-onset preeclampsia and HELLP syndrome. PLoS ONE. 2008 Apr 2; 3 (4): e1865.
- van Wissen S, Bastiaansen BA, Stroobants AK, van den Dool EJ, Idu MM, Levi M, Stoes ES. Catastrophic antiphospholipid syndrome mimicking a malignant pancreatic tumour--a case report. Lupus. 2008; 17 (6): 586-90.
- Vente MA, Nijssen JF, de Wit TC, Seppenwoolde JH, Krijger GC, Seevinck PR, Huisman A, Zonnenberg BA, van den Ingh TS, van het Schip AD. Clinical effects of transcatheter hepatic arterial embolization with holmium-166 poly(L-lactic acid) microspheres in healthy pigs. Eur J Nucl Med Mol Imaging. 2008 Jul; 35 (7): 1259-71.
- Veurink JW, Harteveld CL, Van den Bergh FAJTM. Een Nederlandse familie met zowel beta- als deltathalassemie. Ned Tijdschr Hematol 2008; 5 (7): 290-5.

Analytisch: Immunochemie

- Ballieux BE, Weijl NI, Gelderblom H, van Pelt J, Osanto S. False-positive serum human chorionic gonadotropin (hCG) in a male patient with a malignant germ cell tumor of the testis: A case report and review of the literature. The Oncologist. 2008; 13 (11): 1149-1154
- Boer AK, Weekamp HH, van der Lugt CP, van de Laar MAFJ, van der Sluijs Veer G. De diagnostische accuratesse van de anticitrullinebepaling voor het vaststellen van reumatoïde artritis door de huisarts. Ned Tijdschr Geneesk 2008; 152 (2): 91-5.
- Cobbaert CM, Weykamp CW, Michielsen EC, Baadenhuijsen H, van Diejen-Visser MP. Time-dependent instability of cardiac troponins in human plasma spiked with NIST reference material 2921. Clin Chem. 2008 Dec; 54 (12): 2078-9.

de Graaf AO, de Jonge R, Velmans MH, Swaanenburg JCJM, Janssen MJW. Evaluation of blood collection tubes specific for homocysteine measurement on AxSYM, Immulite and LC-MS/MS systems *Ned Tijdschr Klin Chem Labgeneesk*. 2008; 33 (3): 172-4.

de Graaf AO, de Jonge R, Velmans MH, Swaanenburg JCJM, Janssen MJW. Evaluation of blood collection tubes specific for homocysteine measurement. *Clin Chem Lab Med*. 2008; 46 (11): 1652-4.

Kerkhof D, Peters B, Scharnhorst V. Performance of the Advia Centaur second-generation troponin assay TnI-Ultra compared with the first-generation cTnI assay. *Ann Clin Biochem*. 2008 May; 45 (Pt 3): 316-7.

Tjalsma H, Schaeps RM, Swinkels DW. Immunoproteomics: from biomarker discovery to diagnostic applications. *Proteomics Clin Appl* 2008; 2 (2): 167-80.

van de Kerkhof D, Peters B, Scharnhorst V. Troponin I concentrations in heparinized plasma and serum differ when measured with the Advia Centaur TnI-Ultra assay. *Scand J Clin Lab Invest*. 2008 Apr 11:1-3.

van den Ouweland JM, van Daal H, Klaassen CH, van Aarssen Y, Hartevelde CL, Giordano PC The silent haemoglobin alpha chain variant Hb Riccarton [alpha51(CE9)Gly->Ser] may affect HbA1c determination on the HLC-723 G7 analyzer. *Clin Chem Lab Med*. 2008; 46 (6): 827-30.

van Toorenenbergen AW, Hooijkaas H, Heerenbrink GK, Dufour-van den Goorbergh DM. Heterophilic antibody interference in a tryptase immunoassay. *Clin Biochem*. 2008 Mar; 41 (4-5): 331-4.

Analytisch: Klinische chemie

Abbink FC, Laarman CA, Braam KI, van Wijk JA, Kors WA, Bouman AA, Spreeuwenberg MD, Stoffel-Wagner B, Bokenkamp A. Beta-trace protein is not superior to cystatin C for the estimation of GFR in patients receiving corticosteroids. *Clin Biochem*. 2008 Mar; 41 (4-5): 299-305.

Ackermans MT, Klieverik LP, Endert E, Sauerwein HP, Kalsbeek A, Fliers E. Plasma insulin concentrations during a hyperinsulinaemic clamp: what do we measure? *Ann Clin Biochem*. 2008 Jul; 45 (Pt 4): 429-30.

Arndt T, van der Meijden BB, Wielders JP. Atypical serum transferrin isoform distribution in liver cirrhosis studied by HPLC, capillary electrophoresis and transferrin genotyping. *Clin Chim Acta*. 2008 Aug; 394 (1-2): 42-6.

Bons JA, Wodzig WK, van Dieijen-Visser MP. Protein profiling as a diagnostic tool in clinical chemistry: a review. *Clin Chem Lab Med* 2008; 43 (12): 1281-90.

de Jong WH, Graham KS, de Vries EG, Kema IP. Urinary 5-HIAA measurement using automated on-line solid-phase extraction-high-performance liquid chromatography-tandem mass spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci*. 2008 Jun 1; 868 (1-2): 28-33.

Delanghe JR, Cobbaert C, Galteau MM, Harmoinen A, Jansen R, Kruse R, Laitinen P, Thienpont LM, Wuyts B, Weykamp C, Panteghini M. Trueness verification of actual creatinine assays in the European market demonstrates a disappointing variability that needs substantial improvement. An international study in the framework of the EC4 creatinine standardization working group. *Clin Chem Lab Med*. 2008; 46(9): 1319-25.

Russcher H, van Acker BAC, Boonstra JG, de Rijke YB. Aantonen van chylomicronen in pleuravocht. *Ned Tijdschr Klin Chem Labgeneesk* 2008; 33: 162-165.

Heijboer AC, Otten JH, Jelles WJ, Blankenstein MA, Vroonhof K. Importance of sample size in hospital point-of-care glucose measurements. *Clin Chem Lab Med*. 2008; 46(6): 874-5.

Hempfen C, Wanschers H, van der Sluijs Veer G. A fast liquid chromatographic tandem mass spectrometric method for the simultaneous determination of total homocysteine and methylmalonic acid. *Anal Bioanal Chem*. 2008 May; 391 (1): 263-70.

Sati L, Demir AY, Sarikcioglu L, Demir R. Arrangement of collagen fibers in human placental stem villi. *Acta Histochem*. 2008; 110 (5): 371-9.

Smit NP, van Rossum HH, Romijn FP, Sellar KJ, Breetveld M, Gibbs S, van Pelt J. Calcineurin activity and inhibition in skin and (epi)dermal cell cultures. *J Invest Dermatol*. 2008 Jul; 128 (7): 1686-90.

Vahidnia A, van der Straaten RJ, Romijn F, van Pelt J, van der Voet GB, de Wolff FA. Mechanism of arsenic-induced neurotoxicity may be explained through cleavage of p35 to p25 by calpain. *Toxicol In Vitro*. 2008 Apr; 22 (3): 682-7.

van der Meer IM, Boeke AJ, Lips P, Grootjans-Geerts I, Wuister JD, Devillé WL, Wielders JP, Bouter LM, Middelkoop BJ. Fatty fish and supplements are the greatest modifiable contributors to the serum 25-hydroxyvitamin D concentration in a multiethnic population. *Clin Endocrinol (Oxf)*. 2008 Mar; 68 (3): 466-72.

van Gammeren AJ, van Gool N, de Groot MJM, Cobbaert CM. Analytical performance evaluation of the Cobas 6000 analyzer - special emphasis on trueness verification. *Clin Hem Lab Med*. 2008; 46 (6): 863-71.

van Rossum HH, Romijn FP, Sellar KJ, Smit NP, van der Boog PJ, de Fijter JW, van Pelt J. Variation in leukocyte subset concentrations affects calcineurin activity measurement: implications for pharmacodynamic monitoring strategies. *Clin Chem*. 2008 Mar; 54 (3): 517-24.

van Toorenenbergen AW. Specificity of MMP-9 assays. *J Renin Angiotensin Aldosterone Syst*. 2008 Jun; 9 (2): 65.

Analytisch: Moleculaire technieken

de Waard H, Sonneveld E, de Wit J, Esvelde-van Lange R, Hoeijmakers JH, Vrieling H, van der Horst GT. Cell-type-specific consequences of nucleotide excision repair deficiencies: Embryonic stem cells versus fibroblasts. *DNA Repair (Amst)*. 2008 Oct 1; 7 (10): 1659-69.

Kartikasari AE, Roelofs R, Schaeps RM, Kemna EH, Peters WH, Swinkels DW, Tjalsma H. Secretion of bioactive hepcidin-25 by liver cells correlates with its gene transcription and points towards synergism between iron and inflammation signaling pathways. *Biochim Biophys Acta*. 2008 Dec; 1784 (12): 2029-37.

Neels OC, Koopmans KP, Jager PL, Vercauteren L, van Waarde A, Doorduyn J, Timmer-Bosscha H, Brouwers AH, de Vries EG, Dierckx RA, Kema IP, Elsinga PH. Manipulation of [11C]-5-hydroxytryptophan and 6-[18F]fluoro-3,4-dihydroxy-L-phenylalanine accumulation in neuroendocrine tumor cells. *Cancer Res*. 2008 Sep 1; 68 (17): 7183-90.

Prins J, van der Meijden BB, Kraaijenhagen RJ, Wielders JPM. Inherited chronic obstructive pulmonary disease: new selective-sequencing work-up for α 1-antitrypsin deficiency identifies 2 previously unidentified null alleles. *Clin Chem*. 2008; 54: 101-7.

Ramakers IH, Visser PJ, Aalten P, Bekers O, Slegers K, van Broeckhoven CL, Jolles J, Verhey FR. The association between APOE genotype and memory dysfunction in subjects with mild cognitive impairment is related to age and Alzheimer pathology. *Dement Geriatr Cogn Disord*. 2008; 26 (2): 101-8.

Strunk ALM, Abbes AP, Engel H Real Time PCR-kwantificering van fusiegenen en de detectie van 'minimal-residual disease' op de LightCycler 480 en ABI 7500 Fast. *Ned Tijdschr Klin Chem Labgeneesk*. 2008; 33: 206-8.

Strunk ALM, Joosten H, Abbes AP, Beukhof JR, Engel H. Moleculaire diagnostiek als hulpmiddel bij erfelijke nierziekten. *Ned Tijdschr Klin Chem Labgeneesk*. 2008; 33: 208-10.

Swinkels DW, Girelli D, Laarakkers C, Kroot J, Campostrini N, Kemna EH, Tjalsma H. Advances in quantitative hepcidin measurements by time-of-flight mass spectrometry. *PLoS ONE*. 2008 Jul 16; 3 (7): e2706.

van Schaik RHN, Grandia L, de Goede A, Bet PM, Bekers O, Guchelaar H-J, Goldschmidt HMJ, Meijer MMC, Mulder H, van der Weide J, Deneer VHM. Nederlandse consensus: CYP2D6 genotype '1-0' ingedeeld als intermediaire metaboli-seerder. *Ned Tijdschr Klin Chem Labgeneesk*. 2008; 33 (1): 52-3.

Wijnen PA, Op den Buijsch RA, Cheung SC, van der Heijden J, Hoogtanders K, Stolk LM, van Dieijen-Visser MP, Neef C, Drent M, Bekers O. Genotyping with a dried blood spot method: a useful technique for application in pharmacogenetics. *Clin Chim Acta*. 2008 Feb; 388 (1-2): 189-91.

Bedrijfsvoering: Diversen

Ralser M, Wamelink MM, Struys EA, Joppich C, Krobitsch S, Jakobs C, Lehrach H. A catabolic block does not sufficiently explain how 2-deoxy-D-glucose inhibits cell growth. *Proc Natl Acad Sci U S A*. 2008 Nov 18;105 (46): 17807-11.

Verhaagen J, Janssens PMW. Een ziekenhuislaboratorium is ge-woon heel anders. *Analyse*, nov. 2008: 265-8.

Klinisch: Allergie / infectieziekten

Menheere PPCA. Rapportage van specifieke IgE-bepalingen: voortaan in concentratie. *Ned Tijdschr Klin Chem Labgeneesk*. 2008; 33: 2-6.

Linssen CF, Bekers O, Drent M, Jacobs JA. C-reactive protein and procalcitonin concentrations in bronchoalveolar lavage fluid as a predictor of ventilator-associated pneumonia. *Ann Clin Biochem*. 2008 May; 45 (Pt 3): 293-8.

Robroeks CM, Rosias PP, van Vliet D, Jobsis Q, Yntema JB, Brackel HJ, Damoiseaux JG, den Hartog GM, Wodzig WK, Dompeling E. Biomarkers in exhaled breath condensate indicate presence and severity of cystic fibrosis in children. *Pediatr Allergy Immunol*. 2008 Nov; 19 (7): 652-9.

Robroeks CM, van de Kant KD, van Vliet D, Kester AD, Hendriks HJ, Damoiseaux JG, Wodzig WK, Rijkers GT, Dompeling E, Jobsis Q. Comparison of the anti-inflammatory effects of extra-fine hydrofluoroalkane-beclomethasone vs fluticasone dry powder inhaler on exhaled inflammatory markers in childhood asthma. *Ann Allergy Asthma Immunol*. 2008 Jun; 100 (6): 601-7.

Rosias PP, Robroeks CM, Kester A, den Hartog GJ, Wodzig WK, Rijkers GT, Zimmermann LJ, van Schayck CP, Jobsis Q, Dompeling E. Biomarker reproducibility in exhaled breath condensate collected with different condensers. *Eur Respir J*. 2008 May; 31 (5): 934-42.

Schrama AJ, de Beaufort AJ, Poorthuis BJ, Berger HM, Walther FJ. Secretory phospholipase A(2) in newborn infants with sepsis. *J Perinatol*. 2008 Apr; 28 (4): 291-6.

Tjalsma H, Lambooy L, Hermans PW, Swinkels DW. Shedding & shaving: disclosure of proteomic expressions on a bacterial face. *Proteomics*. 2008 Apr; 8 (7): 1415-28.

Klinisch: Endocrinologie

Albers EM, Riksen-Walraven JM, Sweep FC, de Weerth C. Maternal behavior predicts infant cortisol recovery from a mild everyday stressor. *J Child Psychol Psychiatry*. 2008 Jan; 49 (1): 97-103.

Biermann JC, Holzschleiter L, Kotsch M, Luther T, Kiechle-Bahat M, Sweep FC, Span PN, Schmitt M, Magdolen V. Quantitative RT-PCR assays for the determination of urokinase-type plasminogen activator and plasminogen activator inhibitor type 1 mRNA in primary tumor tissue of breast cancer patients: comparison to antigen quantification by ELISA. *Int J Mol Med*. 2008 Feb; 21 (2): 251-9.

Blumer RM, van der Crabben SN, Stegenga ME, Tanck MW, Ackermans MT, Endert E, van der Poll T, Sauerwein HP. Hyperglycemia prevents the suppressive effect of hyperinsulinemia on plasma adiponectin levels in healthy humans. *Am J Physiol Endocrinol Metab*. 2008 Sep; 295 (3): E613-7.

Claahsen-van der Grinten HL, Otten BJ, Hermus AR, Sweep FC, Hulsbergen-van de Kaa CA. Testicular adrenal rest tumors in patients with congenital adrenal hyperplasia can cause severe testicular damage. *Fertil Steril*. 2008 Mar; 89 (3): 597-601.

Corpeleijn E, Pelsers MM, Soenen S, Mensink M, Bouwman FG, Kooi ME, Saris WH, Glatz JF, Blaak EE. Insulin acutely up-regulates protein expression of the fatty acid transporter CD36 in human skeletal muscle in vivo. *J Physiol Pharmacol*. 2008 Mar; 59 (1): 77-83.

de Kloet CS, Vermetten E, Geuze E, Lentjes EG, Heijnen CJ, Stalla GK, Westenberg HG. Elevated plasma corticotrophin-releasing hormone levels in veterans with posttraumatic stress disorder. *Prog Brain Res*. 2008; 167: 287-91.

de Kort SW, van Dijk M, Willemsen RH, Ester WA, Viet L, de Rijke YB, Hokken-Koelega AC. Cardiovascular risk factors in parents of short children born small for gestational age (SGA). *Pediatr Res*. 2008 Jul; 64 (1): 91-6.

de Lind van Wijngaarden RF, Otten BJ, Festen DA, Joosten KF, de Jong FH, Sweep FC, Hokken-Koelega AC. High prevalence of central adrenal insufficiency in patients with Prader-Willi syndrome. *J Clin Endocrinol Metab*. 2008 May; 93 (5): 1649-54.

de Mast Q, Sweep FC, McCall M, Geurts-Moespot A, Hermesen C, Calandra T, Netea MG, Sauerwein RW, van der Ven AJ. A decrease of plasma macrophage migration inhibitory factor concentration is associated with lower numbers of circulating lymphocytes in experimental *Plasmodium falciparum* malaria. *Parasite Immunol*. 2008 Mar; 30 (3): 133-8.

Freriks K, den Heijer M, Bonenkamp JJ, Biert J, Sweep CG, Hermus AR. Intraoperative parathyroid hormone measurement in patients with primary hyperparathyroidism; particularly valuable for suspected solitary parathyroid adenoma and reoperation. *Ned Tijdschr Geneesk*. 2008 Jan 5; 152 (1): 28-32.

Gilles R, den Heijer M, Ross AH, Sweep FC, Hermus AR, Wet-zels JF. Thyroid function in patients with proteinuria. *Neth J Med*. 2008 Nov; 66 (11): 483-485.

Kok P, Roelfsema F, Frolich M, van Pelt J, Meinders AE, Pijl H. Short-term treatment with bromocriptine improves impaired circadian growth hormone secretion in obese premenopausal women. *J Clin Endocrinol Metab*. 2008 Sep; 93 (9): 3455-61.

Manders RJ, Koopman R, Beelen M, Gijzen AP, Wodzig WK, Saris WH, van Loon LJ. The muscle protein synthetic response to carbohydrate and protein ingestion is not impaired in men with longstanding type 2 diabetes. *J Nutr*. 2008 Jun; 138 (6): 1079-85.

Pelsers MM, Stellingwerff T, van Loon LJ. The role of membrane fatty-acid transporters in regulating skeletal muscle substrate use during exercise. *Sports Med*. 2008; 38 (5): 387-99.

Ross HA, Netea-Maier RT, Schakenraad E, Bravenboer B, Hermus AR, Sweep FC. Assay bias may invalidate decision limits and affect comparability of serum thyroglobulin assay methods: an approach to reduce interpretation differences. *Clin Chim Acta*. 2008 Aug; 394 (1-2): 104-9.

- Schrohl AS, Wurtz S, Kohn E, Banks RE, Nielsen HJ, Sweep FC, Brunner N. Banking of biological fluids for studies of disease-associated protein biomarkers. *Mol Cell Proteomics*. 2008 Oct; 7 (10): 2061-6.
- Strieder TG, Tijssen JG, Wenzel BE, Endert E, Wiersinga WM. Prediction of progression to overt hypothyroidism or hyperthyroidism in female relatives of patients with autoimmune thyroid disease using the Thyroid Events Amsterdam (THEA) score. *Arch Intern Med*. 2008 Aug 11; 168 (15): 1657-63.
- Tanke MA, Kema IP, Dijk-Brouwer J, Doornbos B, de Vries EG, Korf J. Low plasma tryptophan in carcinoid patients is associated with increased urinary cortisol excretion. *Psycho-neuroendocrinology*. 2008 Oct; 33 (9): 1297-301.
- Tran TC, van 't Hof G, Kazemier G, Hop WC, Pek C, van Toorenenbergen AW, van Dekken H, van Eijck CH. Pancreatic fibrosis correlates with exocrine pancreatic insufficiency after pancreatoduodenectomy. *Dig Surg*. 2008 Sep 26; 25 (4): 311-318
- van den Akker EL, Koper JW, van Rossum EF, Dekker MJ, Russcher H, de Jong FH, Uitterlinden AG, Hofman A, Pols HA, Witteman JC, Lamberts SW. Glucocorticoid receptor gene and risk of cardiovascular disease. *Arch Intern Med*. 2008 Jan 14; 168 (1): 33-9.
- van den Bout-van den Beukel CJ, Fievez L, Michels M, Sweep FC, Hermus AR, Bosch ME, Burger DM, Bravenboer B, Koopmans PP, van der Ven AJ. Vitamin D deficiency among HIV Type 1-infected individuals in the Netherlands: effects of antiretroviral therapy. *AIDS Res Hum Retroviruses*. 2008 Nov; 24 (11): 1375-82.
- van der Crabben SN, Allick G, Ackermans MT, Endert E, Romijn JA, Sauerwein HP. Prolonged fasting induces peripheral insulin resistance, which is not ameliorated by high-dose salicylate. *J Clin Endocrinol Metab*. 2008 Feb; 93 (2): 638-41.
- van der Deure WM, Friesema EC, de Jong FJ, de Rijke YB, de Jong FH, Uitterlinden AG, Breteler MM, Peeters RP, Visser TJ. Organic anion transporter 1B1: an important factor in hepatic thyroid hormone and estrogen transport and metabolism. *Endocrinology*. 2008 Sep; 149 (9): 4695-701.
- Vos XG, Smit N, Endert E, Tijssen JG, Wiersinga WM. Frequency and characteristics of TBII-seronegative patients in a population with untreated Graves' hyperthyroidism: a prospective study. *Clin Endocrinol (Oxf)*. 2008 Aug; 69 (2): 311-7.
- Willemsen RH, Mulder PG, van Toorenenbergen AW, Hokken-Koelega AC. Long-term GH treatment is not associated with disadvantageous changes of inflammatory markers and adipocytokines in children born small for gestational age. *Clin Endocrinol (Oxf)*. 2008 Feb; 68 (2): 198-205.
- Willemsen RH, van Dijk M, de Kort SW, van Toorenenbergen AW, Hokken-Koelega AC. Plasma matrix metalloproteinase-9 levels and blood pressure in short children born small for gestational age and effects of growth hormone treatment. *Clin Endocrinol (Oxf)*. 2008 Aug; 69 (2): 264-8.
- Klinisch: Erfelijke stofwisselingsziekten**
- Aerts JM, Groener JE, Kuiper S, Donker-Koopman WE, Strijland A, Ottenhoff R, van Roomen C, Mirzaian M, Wijburg FA, Linthorst GE, Vedder AC, Rombach SM, Cox-Brinkman J, Somerharju P, Boot RG, Hollak CE, Brady RO, Poorthuis BJ. Elevated globotriaosylsphingosine is a hallmark of Fabry disease. *Proc Natl Acad Sci U S A*. 2008 Feb 26; 105 (8): 2812-7.
- Betsalel OT, van de Kamp JM, Martinez-Munoz C, Rosenberg EH, de Brouwer AP, Pouwels PJ, van der Knaap MS, Mancini GM, Jakobs C, Hamel BC, Salomons GS. Detection of low-level somatic and germline mosaicism by denaturing high-performance liquid chromatography in a EURO-MRX family with SLC6A8 deficiency. *Neurogenetics*. 2008 Jul; 9 (3): 183-90.
- Bons JA, Michielsen EC, de Boer D, Bouwman FG, Jaeken J, van Diejen-Visser MP, Rubio-Gozalbo ME, Wodzig WK. A specific immunoprecipitation method for isolating isoforms of insulin-like growth factor binding protein-3 from serum. *Clin Chim Acta*. 2008 Jan; 387 (1-2): 59-65.
- Carrozzo R, Bellini C, Luciola S, Deodato F, Cassandrini D, Cassanello M, Caruso U, Rizzo C, Rizza T, Napolitano ML, Wanders RJA, Jakobs C, Bruno C, Santorelli FM, Dionisi-Vici C, Bonioli E. Peroxisomal acyl-CoA-oxidase deficiency: two new cases. *Am J Med Genet A*. 2008 Jul 1; 146A (13): 1676-81.
- De Feyter HM, Lenaers E, Houten SM, Schrauwen P, Hesselink MK, Wanders RJA, Nicolay K, Prompers JJ. Increased intramyocellular lipid content but normal skeletal muscle mitochondrial oxidative capacity throughout the pathogenesis of type 2 diabetes. *FASEB J*. 2008 Nov; 22 (11): 3947-55.
- Debray FG, Boulanger Y, Khiat A, Decarie JC, Orquin J, Roy MS, Lortie A, Ramos F, Verhoeven NM, Struys E, Blom HJ, Jakobs C, Levy E, Mitchell GA, Lambert M. Reduced brain choline in homocystinuria due to remethylation defects. *Neurology*. 2008 Jul 1; 71 (1): 44-9.
- Derks TG, Boer TS, van Assen A, Bos T, Ruiter J, Waterham HR, Niezen-Koning KE, Wanders RJ, Rondeel JM, Loeber JG, Ten Kate LP, Smit GP, Reijngoud DJ. Neonatal screening for medium-chain acyl-CoA dehydrogenase (MCAD) deficiency in The Netherlands: the importance of enzyme analysis to ascertain true MCAD deficiency. *J Inherit Metab Dis*. 2008 Feb; 31 (1): 88-96.
- Engelen M, Ofman R, Mooijer PA, Poll-The BT, Wanders RJA, Kemp S. Cholesterol-deprivation increases mono-unsaturated very long-chain fatty acids in skin fibroblasts from patients with X-linked adrenoleukodystrophy. *Biochim Biophys Acta*. 2008 Mar; 1781 (3): 105-11.
- van den Bergh FAJTM, Klooster PM, ten Bos R, Straathof-Galema L. 5-Oxoprolinurie: een verworven stofwisselingsstoornis door behandeling met flucloxacilline en paracetamol. *Ned Tijdschr Klin Chem Labgeneesk*. 2008; 33: 39-42.
- Ferdinandusse S, Zomer AW, Komen JC, van den Brink CE, Thanos M, Hamers FP, Wanders RJA, van der Saag PT, Poll-The BT, Brites P. Ataxia with loss of Purkinje cells in a mouse model for Refsum disease. *Proc Natl Acad Sci U S A*. 2008 Nov 18; 105 (46): 17712-7.
- Fons C, Sempere A, Arias A, Lopez-Sala A, Poo P, Pineda M, Mas A, Vilaseca MA, Salomons GS, Ribes A, Artuch R, Campistol J. Arginine supplementation in four patients with X-linked creatine transporter defect. *J Inherit Metab Dis*. 2008 Dec; 31 (6): 724-8.
- Gonzalez F, Schug ZT, Houtkooper RH, MacKenzie ED, Brooks DG, Wanders RJ, Petit PX, Vaz FM, Gottlieb E. Cardiolipin provides an essential activating platform for caspase-8 on mitochondria. *J Cell Biol*. 2008 Nov 17; 183 (4): 681-96.
- Groener JE, Poorthuis BJ, Kuiper S, Hollak CE, Aerts JM. Plasma glucosylceramide and ceramide in type 1 Gaucher disease patients: correlations with disease severity and response to therapeutic intervention. *Biochim Biophys Acta*. 2008 Jan-Feb; 1781 (1-2): 72-8.
- Houtkooper RH, Vaz FM. Cardiolipin, the heart of mitochondrial metabolism. *Cell Mol Life Sci*. 2008 Aug; 65 (16): 2493-506.
- Illsinger S, Lucke T, Peter M, Ruiter JP, Wanders RJA, Deschauer M, Handig I, Wuyts W, Das AM. Carnitine-palmitoyltransferase 2 deficiency: novel mutations and relevance of newborn screening. *Am J Med Genet A*. 2008 Nov 15; 146A (22): 2925-8.

Kornak U, Reynders E, Dimopoulou A, van Reeuwijk J, Fischer B, Rajab A, Budde B, Nurnberg P, Foulquier F; ARCL Debre-type Study Group; Lefeber D, Urban Z, Gruenewald S, Annaert W, Brunner HG, van Bokhoven H, Wevers R, Morava E, Matthijs G, Van Maldergem L, Mundlos S. Impaired glycosylation and cutis laxa caused by mutations in the vesicular H⁺-ATPase subunit ATP6V0A2. *Nat Genet.* 2008 Jan; 40 (1): 32-4.

Kruit JK, Drayer AL, Bloks VW, Blom N, Olthof SG, Sauer PJ, de Haan G, Kema IP, Vellenga E, Kuipers F. Plant sterols cause macrothrombocytopenia in a mouse model of sitosterolemia. *J Biol Chem.* 2008 Mar 7; 283 (10): 6281-7.

Martinez-Munoz C, Rosenberg EH, Jakobs C, Salomons GS. Identification, characterization and cloning of SLC6A8C, a novel splice variant of the creatine transporter gene. *Gene.* 2008 Jul 15; 418 (1-2): 53-9.

Ruijter GJ, Valstar MJ, van de Kamp JM, van der Helm RM, Durand S, van Diggelen OP, Wevers RA, Poorthuis BJ, Pshzhetsky AV, Wijburg FA. Clinical and genetic spectrum of Sanfilippo type C (MPS IIIC) disease in The Netherlands. *Mol Genet Metab.* 2008 Feb; 93 (2): 104-11.

Sanders RJ, Ofman R, Dacremont G, Wanders RJ, Kemp S. Characterization of the human omega-oxidation pathway for omega-hydroxy-very-long-chain fatty acids. *FASEB J.* 2008 Jun; 22 (6): 2064-2071.

van Dijk BA, Laarakkers CM, Klaver SM, Jacobs EM, van Tits LJ, Janssen MC, Swinkels DW. Serum hepcidin levels are innately low in HFE-related haemochromatosis but differ between C282Y-homozygotes with elevated and normal ferritin levels. *Br J Haematol.* 2008 Sep; 142 (6): 979-85.

van Maldegem BT, Duran M, Wanders RJA, Niezen-Koning KE, Hogeveen M, Ijlst L, Waterham HR, Wijburg FA. Short-chain acyl-CoA dehydrogenase deficiency (SCADD): relatively high prevalence in the Netherlands and strongly variable phenotype; neonatal screening not indicated. *Ned Tijdschr Geneesk.* 2008 Jul 26; 152 (30): 1678-85.

van Roermund CW, Visser WF, Ijlst L, van Cruchten A, Boek M, Kulik W, Waterham HR, Wanders RJA. The human peroxisomal ABC half transporter ALDP functions as a homodimer and accepts acyl-CoA esters. *FASEB J.* 2008 Dec; 22 (12): 4201-8.

Wamelink MM, Struys EA, Jakobs C. The biochemistry, metabolism and inherited defects of the pentose phosphate pathway: a review. *J Inher Metab Dis.* 2008 Dec; 31 (6): 703-17.

Wamelink MM, Struys EA, Jansen EE, Levtchenko EN, Zijlstra FS, Engelke U, Blom HJ, Jakobs C, Wevers RA. Sedoheptulokinase deficiency due to a 57-kb deletion in cystinosis patients causes urinary accumulation of sedoheptulose: elucidation of the CARKL gene. *Hum Mutat.* 2008 Apr; 29 (4): 532-6.

Wamelink MM, Struys EA, Salomons GS, Fowler D, Jakobs C, Clayton PT. Transaldolase deficiency in a two-year-old boy with cirrhosis. *Mol Genet Metab.* 2008 Jun; 94 (2): 255-8.

Westermann CM, Dorland L, Votion DM, de Sain-van der Velden MG, Wijnberg ID, Wanders RJA, Spliet WG, Testerink N, Berger R, Ruiten JP, van der Kolk JH. Acquired multiple Acyl-CoA dehydrogenase deficiency in 10 horses with atypical myopathy. *Neuromuscul Disord.* 2008 May; 18 (5): 355-64.

Willemsen AM, Jansen GA, Komen JC, van Hooff S, Waterham HR, Brites PM, Wanders RJA, van Kampen AH. Organization and integration of biomedical knowledge with concept maps for key peroxisomal pathways. *Bioinformatics.* 2008 Aug 15; 24 (16): i21-7.

Zeharia A, Shaag A, Houtkooper RH, Hindi T, de Lonlay P, Erez G, Hubert L, Saada A, de Keyzer Y, Eshel G, Vaz FM, Pines O, Elpeleg O. Mutations in LPIN1 cause recurrent acute myoglobinuria in childhood. *Am J Hum Genet.* 2008 Oct; 83 (4): 489-94.

Klinisch: Gynaecologie / obstetrie / perinatologie

Boxmeer JC, Steegers-Theunissen RP, Lindemans J, Wildhagen MF, Martini E, Steegers EA, Macklon NS. Homocysteine metabolism in the pre-ovulatory follicle during ovarian stimulation. *Hum Reprod.* 2008 Nov; 23 (11): 2570-6.

Brouns R, Ursem N, Lindemans J, Hop W, Pluijm S, Steegers E, Steegers-Theunissen R. Polymorphisms in genes related to folate and cobalamin metabolism and the associations with complex birth defects. *Prenat Diagn.* 2008 Jun; 28 (6): 485-93.

Richir MC, van Leeuwen PA, van den Berg A, Wessels R, Twisk JW, Rauwerda JA, Teerlink T, de Vries TP, van Elburg RM. Plasma ADMA concentrations at birth and mechanical ventilation in preterm infants: a prospective pilot study. *Pediatr Pulmonol.* 2008 Dec; 43 (12): 1161-6.

Verkleij-Hagoort AC, van Driel LM, Lindemans J, Isaacs A, Steegers EA, Helbing WA, Uitterlinden AG, Steegers-Theunissen RP. Genetic and lifestyle factors related to the periconception vitamin B12 status and congenital heart defects: a Dutch case-control study. *Mol Genet Metab.* 2008 May; 94 (1): 112-9.

Klinisch: Hart- en vaatziekten

Aarnoudse AJ, Dieleman JP, Visser LE, Arp PP, van der Heiden IP, van Schaik RH, Molokhia M, Hofman A, Uitterlinden AG, Stricker BH. Common ATP-binding cassette B1 variants are associated with increased digoxin serum concentration. *Pharmacogenet Genomics.* 2008 Apr; 18 (4): 299-305.

Alizadeh Dehnavi R, de Roos A, Rabelink TJ, van Pelt J, Wensink MJ, Romijn JA, Tamsma JT. Elevated CRP levels are associated with increased carotid atherosclerosis independent of visceral obesity. *Atherosclerosis.* 2008 Oct; 200 (2): 417-23.

Alssema M, Schindhelm RK, Dekker JM, Diamant M, Kostense PJ, Teerlink T, Scheffer PG, Nijpels G, Heine RJ. Postprandial glucose and not triglyceride concentrations are associated with carotid intima media thickness in women with normal glucose metabolism: the Hoorn prandial study. *Atherosclerosis.* 2008 Feb; 196 (2): 712-9.

Alssema M, Schindhelm RK, Dekker JM, Diamant M, Nijpels G, Teerlink T, Scheffer PG, Kostense PJ, Heine RJ. Determinants of postprandial triglyceride and glucose responses after two consecutive fat-rich or carbohydrate-rich meals in normoglycemic women and in women with type 2 diabetes mellitus: the Hoorn Prandial Study. *Metabolism.* 2008 Sep; 57 (9): 1262-9.

Bilgin YM, Brand A, Berger SP, Daha MR, Roos A. Mannose-binding lectin is involved in multiple organ dysfunction syndrome after cardiac surgery: effects of blood transfusions. *Transfusion.* 2008 Apr; 48 (4): 601-8.

Bouman HJ, van Werkum JW, Hackeng CM, Verheugt FW, Ten Berg JM. The importance of anticoagulant agents in measuring platelet aggregation in patients treated with clopidogrel and aspirin. *J Thromb Haemost.* 2008 Jun; 6 (6): 1040-2.

Brouwers O, Teerlink T, van Bezu J, Barto R, Stehouwer CD, Schalkwijk CG. Methylglyoxal and methylglyoxal-arginine adducts do not directly inhibit endothelial nitric oxide synthase. *Ann N Y Acad Sci.* 2008 Apr; 1126: 231-4.

Carlstrom M, Brown RD, Edlund J, Sallstrom J, Larsson E, Teerlink T, Palm F, Wahlin N, Persson AE. Role of nitric oxide deficiency in the development of hypertension in hydronephrotic animals. *Am J Physiol Renal Physiol.* 2008 Feb; 294 (2): F362-70.

Crispi F, Hernandez-Andrade E, Pelsers MM, Plasencia W, Benavides-Serralde JA, Eixarch E, Le Noble F, Ahmed A, Glatz JF, Nicolaidis KH, Gratacos E. Cardiac dysfunction and cell damage across clinical stages of severity in growth-restricted fetuses. *Am J Obstet Gynecol.* 2008 Sep; 199 (3): 254.e1-254.e8.

- Ellger B, Langouche L, Richir M, Debaveye Y, Vanhorebeek I, Teerlink T, van Leeuwen PA, van den Berghe G. Modulation of regional nitric oxide metabolism: blood glucose control or insulin? *Intensive Care Med.* 2008 Aug; 34 (8): 1525-33.
- Ellger B, Richir MC, van Leeuwen PA, Debaveye Y, Langouche L, Vanhorebeek I, Teerlink T, Van den Berghe G. Glycemic control modulates arginine and asymmetrical-dimethylarginine levels during critical illness by preserving dimethylarginine-dimethylaminohydrolase activity. *Endocrinology.* 2008 Jun; 149 (6): 3148-57.
- Goei D, Schouten O, Boersma E, Welten GM, Dunkelgrun M, Lindemans J, van Gestel YR, Hoeks SE, Bax JJ, Poldermans D. Influence of renal function on the usefulness of N-terminal pro-B-type natriuretic peptide as a prognostic cardiac risk marker in patients undergoing noncardiac vascular surgery. *Am J Cardiol.* 2008 Jan 1; 101 (1): 122-6.
- Heestermans AA, van Werkum JW, Taubert D, Seesing TH, von Beckerath N, Hackeng CM, Schomig E, Verheugt FW, Ten Berg JM. Impaired bioavailability of clopidogrel in patients with a ST-segment elevation myocardial infarction. *Thromb Res.* 2008; 122 (6): 776-81.
- Hessel MH, Michielsen EC, Atsma DE, Schalijs MJ, van der Valk EJ, Bax WH, Hermens WT, van Diejen-Visser MP, van der Laarse A. Release kinetics of intact and degraded troponin I and T after irreversible cell damage. *Exp Mol Pathol.* 2008 Oct; 85 (2): 90-5.
- IJsselmuiden AJ, Musters RJ, de Ruiter G, van Heerebeek L, Alderse-Baas F, van Schilfgaarde M, Leyte A, Tangelder GJ, Laarman GJ, Paulus WJ. Circulating white blood cells and platelets amplify oxidative stress in heart failure. *Nat Clin Pract Cardiovasc Med.* 2008 Dec; 5 (12): 811-20.
- Lajer M, Tarnow L, Jorsal A, Teerlink T, Parving HH, Rossing P. Plasma concentration of asymmetric dimethylarginine (ADMA) predicts cardiovascular morbidity and mortality in type 1 diabetic patients with diabetic nephropathy. *Diabetes Care.* 2008 Apr; 31 (4): 747-52.
- Landburg PP, Teerlink T, Muskiet FA, Duits AJ, Schnog JJ; CURAMA study group. Plasma concentrations of asymmetric dimethylarginine, an endogenous nitric oxide synthase inhibitor, are elevated in sickle cell patients but do not increase further during painful crisis. *Am J Hematol.* 2008 Jul; 83 (7): 577-9.
- Landburg PP, Teerlink T, van Beers EJ, Muskiet FA, Kappers-Klunne MC, van Esser JW, Mac Gillavry MR, Biemond BJ, Brandjes DP, Duits AJ, Schnog JJ; CURAMA study group. Association of asymmetric dimethylarginine with sickle cell disease-related pulmonary hypertension. *Haematologica.* 2008 Sep; 93 (9): 1410-2.
- Luiken JJ, Vertommen D, Coort SL, Habets DD, El Hasnaoui M, Pelsers MM, Viollet B, Bonen A, Hue L, Rider MH, Glatz JF. Identification of protein kinase D as a novel contraction-activated kinase linked to GLUT4-mediated glucose uptake, independent of AMPK. *Cell Signal.* 2008 Mar; 20 (3): 543-56.
- Lund SS, Tarnow L, Stehouwer CD, Schalkwijk CG, Teerlink T, Gram J, Winther K, Frandsen M, Smidt UM, Pedersen O, Parving HH, Vaag AA. Impact of metformin versus repaglinide on non-glycaemic cardiovascular risk markers related to inflammation and endothelial dysfunction in non-obese patients with type 2 diabetes. *Eur J Endocrinol.* 2008 May; 158 (5): 631-41.
- Michielsen EC, Bakker JA, Kimmenade RR, Pinto YM, Diejen-Visser MP. The diagnostic value of serum and urinary NT-proBNP for heart failure. *Ann Clin Biochem.* 2008 Jul; 45 (Pt 4): 389-94.
- Michielsen EC, Wodzig WK, Van Diejen-Visser MP. Cardiac troponin T release after prolonged strenuous exercise. *Sports Med.* 2008; 38 (5): 425-35.
- Palm F, Friederich M, Carlsson PO, Hansell P, Teerlink T, Liss P. Reduced nitric oxide in diabetic kidneys due to increased hepatic arginine metabolism: implications for renomedullary oxygen availability. *Am J Physiol Renal Physiol.* 2008 Jan; 294 (1): F30-7.
- Richir MC, Bouwman RH, Teerlink T, Siroen MP, de Vries TP, van Leeuwen PA. The prominent role of the liver in the elimination of asymmetric dimethylarginine (ADMA) and the consequences of impaired hepatic function. *J Parenter Enteral Nutr.* 2008 Nov-Dec; 32 (6): 613-21.
- Roest M, Voorbij HAM, van der Schouw YT, Peeters PHM, Teerlink T, Scheffer PG. High levels of urinary F2-isoprostanes predict cardiovascular mortality in postmenopausal women. *J Clin Lipidol.* 2008; 2: 298-303.
- Rutten JH, Steyerberg EW, Boomsma F, van Saase JL, Deckers JW, Hoogsteden HC, Lindemans J, van den Meiracker AH. N-terminal pro-brain natriuretic peptide testing in the emergency department: beneficial effects on hospitalization, costs, and outcome. *Am Heart J.* 2008 Jul; 156 (1): 71-7.
- Scheffer PG, Teerlink T, Dekker JM, Bos G, Nijpels G, Diamant M, Kostense PJ, Stehouwer CD, Heine RJ. Increased plasma apolipoprotein C-III concentration independently predicts cardiovascular mortality: the Hoorn Study. *Clin Chem.* 2008 Aug; 54 (8): 1325-30.
- Schindhelm RK, Alsema M, Diamant M, Teerlink T, Dekker JM, Kok A, Kostense PJ, Nijpels G, Heine RJ, Scheffer PG. Comparison of two consecutive fat-rich and carbohydrate-rich meals on postprandial myeloperoxidase response in women with and without type 2 diabetes mellitus. *Metabolism.* 2008 Feb; 57 (2): 262-7.
- Schindhelm RK, Diamant M. Niet-alcoholische leversteatose: een nieuwe cardiovasculaire risicofactor? *Ned Tijdschr Diabetologie.* 2008; 6 (2): 65-70.
- Siroen MP, Wiest R, Richir MC, Teerlink T, Rauwerda JA, Drescher FT, Zorger N, van Leeuwen PA. Transjugular intrahepatic portosystemic shunt-placement increases arginine/asymmetric dimethylarginine ratio in cirrhotic patients. *World J Gastroenterol.* 2008 Dec 21; 14 (47): 7214-9.
- Teichert M, Visser LE, van Schaik RH, Hofman A, Uitterlinden AG, De Smet PA, Witteman JC, Stricker BH. Vitamin K epoxide reductase complex subunit 1 (VKORC1) polymorphism and aortic calcification: the Rotterdam Study. *Arterioscler Thromb Vasc Biol.* 2008 Apr; 28 (4): 771-6.
- van Driel LM, Smedts HP, Helbing WA, Isaacs A, Lindemans J, Uitterlinden AG, van Duijn CM, de Vries JH, Steegers EA, Steegers-Theunissen RP. Eight-fold increased risk for congenital heart defects in children carrying the nicotinamide N-methyltransferase polymorphism and exposed to medicines and low nicotinamide. *Eur Heart J.* 2008 Jun; 29 (11): 1424-31.
- van Driel LM, Verkleij-Hagoort AC, de Jonge R, Uitterlinden AG, Steegers EA, van Duijn CM, Steegers-Theunissen RP. Two MTHFR polymorphisms, maternal B-vitamin intake, and CHDs. *Birth Defects Res A Clin Mol Teratol.* 2008 Jun; 82 (6): 474-81.
- van Hecke MV, Dekker JM, Nijpels G, Teerlink T, Jakobs C, Stolk RP, Heine RJ, Bouter LM, Polak BC, Stehouwer CD. Homocysteine, S-adenosylmethionine and S-adenosylhomocysteine are associated with retinal microvascular abnormalities: the Hoorn Study. *Clin Sci (Lond).* 2008 Apr; 114 (7): 479-87.
- van Werkum JW, Hackeng CM, Smit JJJ, van 't Hof AWJ, Verheugt FWA, ten Berg JM. Monitoring antiplatelet therapy with point-of-care platelet function assays: a review of the evidence. *Future Cardiol.* 2008; 4 (1): 33-55.
- van Werkum JW, Harmsze AM, Elsenberg EH, Bouman HJ, ten Berg JM, Hackeng CM. The use of the VerifyNow system to monitor antiplatelet therapy: a review of the current evidence. *Platelets.* 2008 Nov; 19 (7): 479-88.

van Werkum JW, Topcu Y, Postma S, Kelder JC, Hackeng CM, ten Berg JM, Verheugt FW. Effects of diabetes mellitus on platelet reactivity after dual antiplatelet therapy with aspirin and clopidogrel. *Thromb Haemost*. 2008 Mar; 99 (3): 637-9.

Klinisch: Hematologie

van Gammeren AJ, Cobbaert CM, Schrauwen CJM, Voets MAJ, Ermens AAM. Fresh frozen plasma transfusion for reversal of prolonged postanesthesia. *Transfus Med*. 2008; 18 (2): 134-6.

Bosman GJ, Werre JM, Willekens FL, Novotny VM. Erythrocyte ageing in vivo and in vitro: structural aspects and implications for transfusion. *Transfus Med*. 2008 Dec; 18 (6): 335-47.

Brouwer HB, Kemna EHJM, Swinkels DW. Hepcidine, een ijzersterke biomarker? *Ned Tijdschr Hematol*. 2008; 5: 102-8.

Brouwers MC, Govers-Riemslog J, Schalkwijk CG, van Greevenbroek MM, van der Kallen CJ, Bekers O, van Dieijen-Visser MP, Ten Oever J, Bilderbeek-Beckers MA, de Bruin TW, Ten Cate H, Stehouwer CD. Plasma PAI-1 levels are independently related to fatty liver and hypertriglyceridemia in familial combined hyperlipidemia, involvement of apolipoprotein E. *Thromb Res*. 2008; 122 (4): 466-72.

Cherian S, Forbes DA, Cook AG, Sanfilippo FM, Kemna EH, Swinkels DW, Burgner DP. An insight into the relationships between hepcidin, anemia, infections and inflammatory cytokines in pediatric refugees: a cross-sectional study. *PLoS ONE*. 2008; 3 (12): e4030.

de Vooght KM, van Wijk R, van Solinge WW. GATA-1 binding sites in exon 1 direct erythroid-specific transcription of PPOX. *Gene*. 2008 Feb 15; 409 (1-2): 83-91.

Engbers-Buijtenhuijs P, Veurink JW, Hartevelde CL, van den Bergh FAJTM, Slomp J. Een Nederlandse familie met zowel beta- als delatthalassemie. *Ned Tijdschr Hematol*. 2008; 5: 290-5.

Ponjee GAE, Khorsand N, Scharnhorst V. Hemostatica. *Ned Tijdschr Klin Chem Labgeneesk*. 2008; 33: 254-9.

Moonen AF, Pilot P, Meijers WG, Waelen RA, Leers MP, Grimm B, Heyligers IC. Filters in autologous blood retransfusion systems affect the amount of blood cells retransfused in total knee arthroplasty: a pilot study. *Acta Orthop Belg*. 2008 Apr; 74 (2): 210-5.

Remijn JA, Lucas S, Wildeboer B, van Suijlen JD, Adriaansen HJ. Strongly increased international normalized ratio with recombinant Neoplastin R compared with tissue extract Neoplastin Plus in patients initiating oral anticoagulant therapy: implications for anticoagulation dosage. *Clin Chem*. 2008 Nov; 54 (11): 1929-31.

Tamary H, Shalev H, Perez-Avraham G, Zoldan M, Levi I, Swinkels DW, Tanno T, Miller JL. Elevated growth differentiation factor 15 expression in patients with congenital dyserythropoietic anemia type I. *Blood*. 2008 Dec 15; 112 (13): 5241-4.

van Gammeren AJ, Overbeeke MAM, Idema RN, van Beek RHT, Ten Kate-Booij MJ, Ermens AAM. Haemolytic disease of the newborn because of rare anti-*Vel*. *Transfus Med*. 2008; 18, 197-8.

Vlasveld LT, de Wit CW, Verweij RA, Castel A, Jansen PM, Peters AA. Myomatous erythrocytosis syndrome: further proof for the pathogenic role of erythropoietin. *Neth J Med*. 2008 Jul-Aug; 66(7): 283-5.

Willekens FL, Werre JM, Groenen-Dopp YA, Roerdinkholder-Stoelwinder B, de Pauw B, Bosman GJ. Erythrocyte vesiculation: a self-protective mechanism? *Br J Haematol*. 2008 May; 141 (4): 549-56.

Klinisch: Interne geneeskunde

Allegaert K, van den Anker JN, de Hoon JN, van Schaik RH, Debeer A, Tibboel D, Naulaers G, Anderson BJ. Covariates of tramadol disposition in the first months of life. *Br J Anaesth*. 2008 Apr; 100 (4): 525-32.

Allegaert K, van Schaik RH, Vermeersch S, Verbesselt R, Cossey V, Vanhole C, van Fessem M, de Hoon J, van den Anker JN. Postmenstrual age and CYP2D6 polymorphisms determine tramadol o-demethylation in critically ill neonates and infants. *Pediatr Res*. 2008 Jun; 63 (6): 674-9.

Bons JAP, van der Heijde D, Bouwman FG, Mariman EC, van Dieijen-Visser MP, Landewe R, van der Linden S, Wodzig WKWH. Searching for potential biomarkers that can be used as biomarkers for diagnosing ankylosing spondylitis. *Biomarkers in Medicine* 2008; 2 (1): 23-30

Darbari DS, van Schaik RH, Capparelli EV, Rana S, McCarter R, van den Anker J. UGT2B7 promoter variant -840G>A contributes to the variability in hepatic clearance of morphine in patients with sickle cell disease. *Am J Hematol*. 2008 Mar; 83 (3): 200-2.

Hesselink DA, van Schaik RH, Nauta J, van Gelder T. A drug transporter for all ages? ABCB1 and the developmental pharmacogenetics of cyclosporine. *Pharmacogenomics*. 2008 Jun; 9 (6): 783-9.

Hesselink DA, van Schaik RH, van Agteren M, de Fijter JW, Hartmann A, Zeier M, Budde K, Kuypers DR, Pisarski P, Meur YL, Mamelok RD, van Gelder T. CYP3A5 genotype is not associated with a higher risk of acute rejection in tacrolimus-treated renal transplant recipients. *Pharmacogenet Genomics*. 2008 Apr; 18 (4): 339-348.

Hunfeld NG, Mathot RA, Touw DJ, van Schaik RH, Mulder PG, Franck PF, Kuipers EJ, Geus WP. Effect of CYP2C19*2 and *17 mutations on pharmacodynamics and kinetics of proton pump inhibitors in Caucasians. *Br J Clin Pharmacol*. 2008 May; 65(5): 752-60.

Kemna EH, Tjalsma H, Willems HL, Swinkels DW. Hepcidin: from discovery to differential diagnosis. *Haematologica*. 2008 Jan; 93 (1): 90-7.

Moues CM, van Toorenenbergen AW, Heule F, Hop WC, Hovius SE. The role of topical negative pressure in wound repair: Expression of biochemical markers in wound fluid during wound healing. *Wound Repair Regen*. 2008 Jul-Aug; 16 (4): 488-94.

Norton L, Parr T, Chokkalingam K, Bardsley RG, Ye H, Bell GI, Pelsers MM, van Loon LJ, Tsintzas K. Calpain-10 gene and protein expression in human skeletal muscle: effect of acute lipid-induced insulin resistance and type 2 diabetes. *J Clin Endocrinol Metab*. 2008 Mar; 93 (3): 992-8.

Oortwijn BD, Eijgenraam JW, Rastaldi MP, Roos A, Daha MR, van Kooten C. The role of secretory IgA and complement in IgA nephropathy. *Semin Nephrol*. 2008 Jan; 28 (1): 58-65.

Pelsers MM. Fatty acid-binding protein as marker for renal injury. *Scand J Clin Lab Invest Suppl*. 2008; 241: 73-7.

Peters JH, Wierdsma NJ, Teerlink T, van Leeuwen PA, Mulder CJ, van Bodegraven AA. The citrulline generation test: proposal for a new enterocyte function test. *Aliment Pharmacol Ther*. 2008 Jun; 27 (12): 1300-10.

Swinkels DW, Wetzels JF. Hepcidin: a new tool in the management of anaemia in patients with chronic kidney disease? *Nephrol Dial Transplant*. 2008 Aug; 23(8):2450-3.

van Agteren M, Armstrong VW, van Schaik RH, de Fijter H, Hartmann A, Zeier M, Budde K, Kuypers D, Pisarski P, Le Meur Y, van der Werf M, Mamelok RD, Oellerich M, van Gelder T. AcylMPAG plasma concentrations and mycophenolic acid-related side effects in patients undergoing renal transplantation are not related to the UGT2B7-840G>A gene polymorphism. *Ther Drug Monit.* 2008 Aug; 30 (4): 439-44.

van Stijn MF, Ligthart-Melis GC, Boelens PG, Scheffer PG, Teerlink T, Twisk JW, Houdijk AP, van Leeuwen PA. Antioxidant enriched enteral nutrition and oxidative stress after major gastrointestinal tract surgery. *World J Gastroenterol.* 2008 Dec 7; 14 (45): 6960-9.

Verschuren JJ, Roos A, Schaapherder AF, Mallat MJ, Daha MR, de Fijter JW, Berger SP. Infectious complications after simultaneous pancreas-kidney transplantation: a role for the lectin pathway of complement activation. *Transplantation.* 2008 Jan 15; 85 (1): 75-80.

Wijnen PA, Drent M, Nelemans PJ, Kuijpers PM, Koek GH, Neef C, Haenen GR, Bekers O. Role of cytochrome p450 polymorphisms in the development of pulmonary drug toxicity: a case-control study in the Netherlands. *Drug Saf.* 2008; 31 (12): 1125-34.

Yazdanpanah N, Uitterlinden AG, Zillikens MC, Jhamai M, Rivadeneira F, Hofman A, de Jonge R, Lindemans J, Pols HA, van Meurs JB. Low Dietary Riboflavin but Not Folate Predicts Increased Fracture Risk in Postmenopausal Women Homozygous for the MTHFR 677 T Allele. *J Bone Miner Res.* 2008 Jan; 23 (1): 86-94.

Klinisch: Neurologie / psychiatrie

Bijl MJ, Visser LE, Hofman A, Vulto AG, van Gelder T, Stricker BH, van Schaik RH. Influence of the CYP2D6*4 polymorphism on dose, switching and discontinuation of antidepressants. *Br J Clin Pharmacol.* 2008 Apr; 65 (4): 558-64.

de Kloet C, Vermetten E, Lentjes E, Geuze E, van Pelt J, Manuel R, Heijnen C, Westenberg H. Differences in the response to the combined DEX-CRH test between PTSD patients with and without co-morbid depressive disorder. *Psychoneuroendocrinology.* 2008 Apr; 33 (3): 313-20.

Elzinga BM, Roelofs K, Tollenaar MS, Bakvis P, J. van Pelt J, Spinhoven P. Diminished cortisol responses to psychosocial stress associated with lifetime adverse events. A study among healthy young subjects. *Psychoneuroendocrinology* 2008; 33 (2): 227-37.

Giltay EJ, Bunck MC, Gooren LJ, Zitman FG, Diamant M, Teerlink T. Effects of sex steroids on the neurotransmitter-specific aromatic amino acids phenylalanine, tyrosine, and tryptophan in transsexual subjects. *Neuroendocrinology.* 2008; 88 (2): 103-10.

Jabbi M, Korf J, Ormel J, Kema IP, den Boer JA. Investigating the molecular basis of major depressive disorder etiology: a functional convergent genetic approach. *Ann N Y Acad Sci.* 2008 Dec; 1148: 42-56.

Miners JS, Verbeek MM, Rikkert MO, Kehoe PG, Love S. Immunocapture-based fluorometric assay for the measurement of neprilysin-specific enzyme activity in brain tissue homogenates and cerebrospinal fluid. *J Neurosci Methods.* 2008 Jan 30; 167 (2): 229-36.

Schenk PW, van Fessem MA, Verploegh-van Rij S, Mathot RA, van Gelder T, Vulto AG, van Vliet M, Lindemans J, Bruijn JA, van Schaik RH. Association of graded allele-specific changes in CYP2D6 function with imipramine dose requirement in a large group of depressed patients. *Mol Psychiatry.* 2008 Jun; 13 (6): 597-605.

Smarius LJ, Jacobs GE, Hoeberechts-Lefrandt DH, de Kam ML, van der Post JP, de Rijk R, van Pelt J, Schoemaker RC,

Zitman FG, van Gerven JM, Gijsman HJ. Pharmacology of rising oral doses of 5-hydroxytryptophan with carbidopa. *J Psychopharmacol.* 2008 Jun; 22 (4): 426-33.

Smolders J, Damoiseaux J, Menheere P, Hupperts R. Vitamin D as an immune modulator in multiple sclerosis, a review. *J Neuroimmunol.* 2008 Feb; 194 (1-2): 7-17.

Staals J, Bons JA, van Oostenbrugge RJ, Knottnerus IL, van Dieijen-Visser MP, Bouwman FG, Mariman EC, Delanghe JR, Lodder J, Wodzig WK. A SELDI-TOF-MS Study in Lacunar Stroke with Subsequent Haptoglobin Phenotyping. *Curr Neurovasc Res.* 2008 May; 5 (2): 93-8.

van Veen JF, van Vliet IM, Derijk RH, van Pelt J, Mertens B, Zitman FG. Elevated alpha-amylase but not cortisol in generalized social anxiety disorder. *Psychoneuroendocrinology.* 2008 Nov; 33 (10): 1313-21.

Verwey NA, Bouwman FH, van der Flier WM, Veerhuis R, Scheltens P, Blankenstein MA. Variability in longitudinal cerebrospinal fluid tau and phosphorylated tau measurements. *Clin Chem Lab Med.* 2008; 46 (9): 1300-4.

Klinisch: Oncologie

Bayer C, Schilling D, Hoetzel J, Egermann HP, Zips D, Yaromina A, Geurts-Moespot A, Sprague LD, Sweep F, Baumann M, Molls M, Adam M. PAI-1 levels predict response to fractionated irradiation in 10 human squamous cell carcinoma lines of the head and neck. *Radiother Oncol.* 2008 Mar; 86 (3): 361-8.

Broeveer FJ, Osanto S, Ritsema van Eck HJ, van Steijn AQ, Balleux BE, Schoemaker RC, Cohen AF, Burggraaf J. Evaluation of biomarkers for cardiotoxicity of anthracyclin-based chemotherapy. *J Cancer Res Clin Oncol.* 2008 Sep; 134 (9): 961-8.

Donker ML, van Doormaal JJ, van Doormaal FF, Kluin PM, van der Veer E, de Monchy JG, Kema IP, Kluin-Nelemans HC. Biochemical markers predictive for bone marrow involvement in systemic mastocytosis. *Haematologica.* 2008 Jan; 93 (1): 120-3.

Eekers DB, Laschet A, de Groot M, Roelofs E, Kester A, De-laere K, Lambin P, van Gils F, Nap M, ten Kate J. Why determine only the total prostate-specific antigen, if the free-to-total ratio contains the information? *Ann Clin Biochem.* 2008 May; 45 (Pt 3): 270-4.

Gudbjartsson DF, Walters GB, Thorleifsson G, Stefansson H, Halldorsson BV, Zusmanovich P, Sulem P, Thorlacius S, Gylfason A, Steinberg S, Helgadóttir A, Ingason A, Steinthorsdóttir V, Olafsdóttir EJ, Olafsdóttir GH, Jonsson T, Borch-Johnsen K, Hansen T, Andersen G, Jorgensen T, Pedersen O, Aben KK, Witjes JA, Swinkels DW, den Heijer M, Franke B, Verbeek AL, Becker DM, Yanek LR, Becker LC, Tryggvadóttir L, Rafnar T, Gulcher J, Kiemeneý LA, Kong A, Thorsteinsdóttir U, Stefansson K. Many sequence variants affecting diversity of adult human height. *Nat Genet.* 2008 May; 40 (5): 609-15.

Gudmundsson J, Sulem P, Rafnar T, Bergthorsson JT, Manolescu A, Gudbjartsson D, Agnarsson BA, Sigurdsson A, Benediksdóttir KR, Blondal T, Jakobsdóttir M, Stacey SN, Kostic J, Kristinsson KT, Birgisdóttir B, Ghosh S, Magnúsdóttir DN, Thorlacius S, Thorleifsson G, Zheng SL, Sun J, Chang BL, Elmore JB, Breyer JP, McReynolds KM, Bradley KM, Yaspan BL, Wiklund F, Stattin P, Lindstrom S, Adami HO, McDonnell SK, Schaid DJ, Cunningham JM, Wang L, Cerhan JR, St Sauver JL, Isaacs SD, Wiley KE, Partin AW, Walsh PC, Polo S, Ruiz-Echarri M, Navarrete S, Fuertes F, Saez B, Godino J, Weijerman PC, Swinkels DW, Aben KK, Witjes JA, Suarez BK, Helfand BT, Frigge ML, Kristjánsson K, Ober C, Jonsson E, Einarsson GV, Xu J, Gronberg H, Smith JR, Thibodeau SN, Isaacs WB, Catalona WJ, Mayordomo JI, Kiemeneý LA, Barkardóttir RB, Gulcher JR, Thorsteinsdóttir U, Kong A, Stefansson K. Common sequence variants on 2p15 and Xp11.22 confer susceptibility to prostate cancer. *Nat Genet.* 2008 Mar; 40 (3): 281-3.

- Gutierrez-Fernandez A, Fueyo A, Folgueras AR, Garabaya C, Pennington CJ, Pilgrim S, Edwards DR, Holliday DL, Jones JL, Span PN, Sweep FC, Puente XS, Lopez-Otin C. Matrix metalloproteinase-8 functions as a metastasis suppressor through modulation of tumor cell adhesion and invasion. *Cancer Res.* 2008 Apr 15; 68 (8): 2755-63.
- Herwig R, Mitteregger D, Djavan B, Kramer G, Margreiter M, Leers MP, Glodny B, Haider DG, Horl WH, Marberger M. Detecting prostate cancer by intracellular macrophage prostate-specific antigen (PSA): a more specific and sensitive marker than conventional serum total PSA. *Eur J Clin Invest.* 2008 Jun; 38 (6): 430-7.
- Huang L, Tissing WJ, de Jonge R, van Zelst BD, Pieters R. Polymorphisms in folate-related genes: association with side effects of high-dose methotrexate in childhood acute lymphoblastic leukemia. *Leukemia.* 2008 Sep; 22 (9): 1798-800.
- Jansen FH, Roobol M, Bangma CH, van Schaik RH. Clinical impact of new prostate-specific antigen WHO standardization on biopsy rates and cancer detection. *Clin Chem.* 2008 Dec; 54 (12): 1999-2006.
- Kiemeny LA, Thorlacius S, Sulem P, Geller F, Aben KK, Stacey SN, Gudmundsson J, Jakobsdottir M, Bergthorsson JT, Sigurdsson A, Blondal T, Witjes JA, Vermeulen SH, Hulsbergen-van de Kaa CA, Swinkels DW, Ploeg M, Cornel EB, Vergunst H, Thorgeirsson TE, Gudbjartsson D, Gudjonsson SA, Thorleifsson G, Kristinsson KT, Mouy M, Snorraddottir S, Placidi D, Campagna M, Arici C, Koppova K, Gurzau E, Rudnai P, Kellen E, Polidoro S, Guarrera S, Sacerdote C, Sanchez M, Saez B, Valdivia G, Ryk C, de Verdier P, Lindblom A, Golka K, Bishop DT, Knowles MA, Nikulasson S, Petursdottir V, Jonsson E, Geirsson G, Kristjansson B, Mayordomo JI, Steineck G, Porru S, Buntinx F, Zeegers MP, Fletcher T, Kumar R, Matullo G, Vineis P, Kiltie AE, Gulcher JR, Thorsteinsdottir U, Kong A, Rafnar T, Stefansson K. Sequence variant on 8q24 confers susceptibility to urinary bladder cancer. *Nat Genet.* 2008 Nov; 40 (11): 1307-12.
- Koopmans KP, de Groot JW, Plukker JT, de Vries EG, Kema IP, Sluiter WJ, Jager PL, Links TP. 18F-dihydroxyphenylalanine PET in patients with biochemical evidence of medullary thyroid cancer: relation to tumor differentiation. *J Nucl Med.* 2008 Apr; 49 (4): 524-31.
- Koopmans KP, Jager PL, Kema IP, Kerstens MN, Albers F, Dullaart RP. 111In-octreotide is superior to 123I-metaiodobenzylguanidine for scintigraphic detection of head and neck paragangliomas. *J Nucl Med.* 2008 Aug; 49 (8): 1232-7.
- Koopmans KP, Neels OC, Kema IP, Elsinga PH, Sluiter WJ, Vanghillewe K, Brouwers AH, Jager PL, de Vries EG. Improved staging of patients with carcinoid and islet cell tumors with 18F-dihydroxy-phenyl-alanine and 11C-5-hydroxy-tryptophan positron emission tomography. *J Clin Oncol.* 2008 Mar 20; 26 (9): 1489-95.
- Leers MP, Nap M, Herwig R, Delaere K, Nauwelaers F. Circulating PSA-containing macrophages as a possible target for the detection of prostate cancer: A three-color/five-parameter flow cytometric Study on peripheral blood samples. *Am J Clin Pathol.* 2008 Apr; 129 (4): 649-56.
- McKiernan E, O'Brien K, Grebenchtchikov N, Geurts-Moespot A, Sieuwerts AM, Martens JW, Magdolen V, Evoy D, McDermott E, Crown J, Sweep FC, Duffy MJ. Protein kinase Cdelta expression in breast cancer as measured by real-time PCR, western blotting and ELISA. *Br J Cancer.* 2008 Nov 18; 99 (10): 1644-50.
- Oei AL, Moreno M, Verheijen RH, Sweep FC, Thomas CM, Massuger LF, von Mensdorff-Pouilly S. Induction of IgG antibodies to MUC1 and survival in patients with epithelial ovarian cancer. *Int J Cancer.* 2008 Oct 15; 123 (8): 1848-53.
- Oei AL, Sweep FC, Geurts-Moespot A, van Tienoven D, Von Mensdorff-Pouilly S, Thomas CM, Massuger LF. Human anti-mouse IgM and IgG responses in ovarian cancer patients after radioimmunotherapy with 90Y-muHMF1. *Anticancer Res.* 2008 Sep-Oct; 28 (5A): 2721-5.
- Oei AL, Sweep FC, Massuger LF, Olthaar AJ, Thomas CM. Transient human anti-mouse antibodies (HAMA) interference in CA 125 measurements during monitoring of ovarian cancer patients treated with murine monoclonal antibody. *Gynecol Oncol.* 2008 May; 109 (2): 199-202.
- Oei AL, Sweep FC, Thomas CM, Boerman OC, Massuger LF. The use of monoclonal antibodies for the treatment of epithelial ovarian cancer (review). *Int J Oncol.* 2008 Jun; 32 (6): 1145-57.
- Scheer MG, Stollman TH, Boerman OC, Verrijp K, Sweep FC, Leenders WP, Ruers TJ, Oyen WJ. Imaging liver metastases of colorectal cancer patients with radiolabelled bevacizumab: Lack of correlation with VEGF-A expression. *Eur J Cancer.* 2008 Sep; 44 (13): 1835-40.
- Smit NP, van Nieuwpoort FA, Marrot L, Out C, Poorthuis B, van Pelt H, Meunier JR, Pavel S. Increased melanogenesis is a risk factor for oxidative DNA damage- Study on cultured melanocytes and atypical nevus cells. *Photochem Photobiol.* 2008 May-Jun; 84 (3): 550-5.
- Span PN, Witjes JA, Grebenchtchikov N, Geurts-Moespot A, Moonen PM, Aalders TW, Vriesema JL, Kiemeny LA, Schalken JA, Sweep FC. Components of the plasminogen activator system and their complexes in renal cell and bladder cancer: comparison between normal and matched cancerous tissues. *BJU Int.* 2008 Jul; 102 (2): 177-82.
- Stacey SN, Manolescu A, Sulem P, Thorlacius S, Gudjonsson SA, Jonsson GF, Jakobsdottir M, Bergthorsson JT, Gudmundsson J, Aben KK, Strobbe LJ, Swinkels DW, van Engelenburg KC, Henderson BE, Kolonel LN, Le Marchand L, Millastre E, Andres R, Saez B, Lambea J, Godino J, Polo E, Tres A, Picelli S, Rantala J, Margolin S, Jonsson T, Sigurdsson H, Jonsdottir T, Hrafnkelsson J, Johannsson J, Sveinsson T, Myrdal G, Grimsson HN, Sveinsdottir SG, Alexiusdottir K, Saemundsdottir J, Sigurdsson A, Kostic J, Gudmundsson L, Kristjansson K, Masson G, Fackenthal JD, Adebamowo C, Ogundiran T, Olopade OI, Haiman CA, Lindblom A, Mayordomo JI, Kiemeny LA, Gulcher JR, Rafnar T, Thorsteinsdottir U, Johannsson OT, Kong A, Stefansson K. Common variants on chromosome 5p12 confer susceptibility to estrogen receptor-positive breast cancer. *Nat Genet.* 2008 Jun; 40 (6): 703-6.
- Thomas CM, Boss EA, Boonstra H, van Tienoven D, Sweep CG, Massuger LF. Gonadotropins and female sex steroid hormones in cyst fluid and serum from patients with ovarian tumors. *Eur J Gynaecol Oncol.* 2008; 29 (5): 468-72.
- van Schaik RH. CYP450 pharmacogenetics for personalizing cancer therapy. *Drug Resist Updat.* 2008 Jun; 11 (3): 77-98.
- van Trommel NE, Ngo Duc H, Massuger LF, Schijf CP, Sweep CG, Thomas CM; Dutch Working Party on Trophoblastic Tumors. Early identification of persistent trophoblastic disease with serum hCG concentration ratios. *Int J Gynecol Cancer.* 2008 Mar-Apr; 18 (2): 318-23.
- Zhong JL, Poghosyan Z, Pennington CJ, Scott X, Handsley MM, Warn A, Gavrilovic J, Honert K, Kruger A, Span PN, Sweep FC, Edwards DR. Distinct functions of natural ADAM-15 cytoplasmic domain variants in human mammary carcinoma. *Mol Cancer Res.* 2008 Mar; 6 (3): 383-94.